

# Measuring Nuclear Effects with MINERvA

APS April Meeting 2011



G. Arturo Fiorentini  
Centro Brasileiro de Pesquisas Físicas

On behalf of the MINERvA collaboration

# MINERvA



## (Main INjector ExpeRiment $\nu$ -A)

- Neutrino scattering experiment in the 1-20 GeV energy range.
- Uses NuMI neutrino beamline at Fermilab.
- Located in front of MINOS near detector.
- Uses MINOS near detector as muon spectrometer.




Fermilab, Batavia IL, USA

# NuMI Beam




- Very intense neutrino beam with a power of 300 – 350 kW and  $\sim 35 \times 10^{12}$  P.O.T. (Protons On Target) per spill.
  - Spill: 10  $\mu$ s duration at  $\sim 0.5$  Hz frequency.
- Energy distribution can be tuned by changing position of target with respect to horns.
- Anti-neutrino beam is obtained by reversing the current in the magnetic horns to focus  $\pi^-$  instead of  $\pi^+$ .

# MINERvA Detector Layout





120 modules of 4 types: nuclear targets, tracker, ECal and HCal.

# MINERvA Modules





=




127 scintillator strips per plane.

Tracker module = 2 planes  
ECAL module = 2 planes + 2 (2 mm thick) sheet of lead  
HCAL module = 1 plane + 1 (1 inch thick) sheet of steel





Triangular strip to allow charge sharing





# MINERvA Nuclear Targets


Liquid Helium



Prototype Water Target




Iron / Lead




Lead / Iron




Carbon / Lead / Iron



Lead



Iron / Lead




# Event Rates

- MINERvA has been approved for  $4.9 \times 10^{20}$  P.O.T. in the low energy neutrino beam
  - So far, we have received  $1.2 \times 10^{20}$  P.O.T. (24% of total)

Expected Event Rates for Low Energy Neutrino-tuned beam		
Target	Fiducial Mass	$\nu_\mu$ CC Events in $1.2 \times 10^{20}$ P.O.T.
Plastic	6.43 tons	409k
Helium	0.25 tons	16.8k
Carbon	0.17 tons	10.8k
Water	0.39 tons	24.4k
Iron	0.97 tons	64.5k
Lead	0.98 tons	68.4k

Some CC inclusive events:




# Nuclear Effects with MINERvA

- Designed to measure neutrino-nucleus cross-section in C, Fe, Pb, He and H<sub>2</sub>O.
  - Having all targets in the same detector and same beam reduces systematic errors in comparisons.
- Study nuclear effects and the A-dependence in neutrino interactions.
- Status on the analysis to compare charge current inclusive cross sections in plastic, Fe and Pb as function of muon energy.





# Nuclear Targets in Event Display

CC event candidate from last target




# Nuclear Targets Event Selection

CC event candidate from last target




# Plastic Reference Target

- All we do for the passive target can be done for a plastic reference target.
  - Compare plastic to Pb and Fe.
  - Minimize systematic errors (efficiency, area, acceptance)




# Muon Energy

$\nu_{\mu}$  CC events




Area-Normalized


Statistical Error Only




Passive Target Event Selection



Active Target Event Selection



# Lead to Iron Ratio



- Ratio removes dependence on neutrino flux.
  - **Lead** and **Iron** see the same beam.


Coming Soon

- Using plastic reference targets minimizes systematic error from different X:Y regions.




# Plastic to Plastic Ratio

- Add another reference target downstream to compare plastic with plastic.
- Measure differences in acceptance between this two positions.



May 2, 2011



APS April Meeting - G. A. Fiorentini

# Summary

- MINERvA is an experiment in the NuMI beamline aiming to improve neutrino cross section measurements and studying nuclear effects.
- Currently working to produce ratios of CC inclusive cross sections of  $\nu_\mu$  on plastic, lead and iron.
- Stay tuned for new results.

# Thanks for listening

## The Minerva Collaboration

G. Tzanakos  
*University of Athens*

J. Cravens, M. Jerkins, S. Kopp, L. Loiacono, J. Ratchford, R. Stevens IV  
*University of Texas at Austin*

D.A.M. Caicedo, C.M. Castromonte, H. da Motta, G. A. Fiorentini, J.L. Palomino  
*Centro Brasileiro de Pesquisas Fisicas*

J. Grange, J. Mousseau, B. Osmanov, H. Ray  
*University of Florida*

D. Boehnlein, R. DeMaat, N. Grossman, D. A. Harris, J. G. Morfn, J. Osta,  
R. B. Pahlka, P. Rubinov, D. W. Schmitz, F.D. Snider, R. Stefanski  
*Fermilab*

J. Felix, A. Higuera, Z. Urrutia, G. Zavala  
*Universidad de Guanajuato*

M.E. Christy, C. Keppel, P. Monagham, T. Walton, L. Y. Zhu  
*Hampton University*

A. Butkevich, S.A. Kulagin  
*Inst. Nucl. Reas. Moscow*

G. Niculescu, I. Niculescu  
*James Madison University*

E. Maher  
*Mass. Col. Lib. Arts*

L. Fields, B. Gobbi, L. Patrick, H. Schellman  
*Northwestern University*

N. Tagg  
*Otterbein College*

S. Boyd, I. Danko, S.A. Dytmann, B. Eberly, Z. Isvan, D. Naples, V. Paolone  
*University of Pittsburgh*

A. M. Gago, N. Ochoa, J.P. Velasquez  
*Pontificia Universidad Catolica del Peru*

S. Avvakumov, A. Bodek, R. Bradford, H. Budd, J. Chvojka, M. Day, H. Lee, S. Manly,  
C. Marshall, K.S. McFarland, A. M. McGowan, A. Mislivec, J. Park, G. Perdue, J. Wolcott  
*University of Rochester*

G. J. Kumbartzki, T. Le, R. D. Ransome, E. C. Schulte, B. G. Tice  
*Rutgers University*

H. Gallagher, T. Kafka, W.A. Mann, W. P. Oliver  
*Tufts University*

C. Simon, B. Ziemer  
*University of California at Irvine*

R. Gran, M. Lanari  
*University of Minnesota at Duluth*

M. Alania, A. Chamorro, K. Hurtado, C. J. Solano Salinas  
*Universidad Nacional de Ingeniera*

W. K. Brooks, E. Carquin, G. Maggi, C. Pea, I.K. Potashnikova, F. Prokoshin  
*Universidad Tecnica Federico Santa Mara*

L. Aliaga, J. Devan, M. Kordosky, J.K. Nelson, J. Walding, D. Zhang  
*College of William and Mary*

# Backups

# MINERvA objectives

- Provide more precise charged current quasi-elastic cross sections to oscillation experiments
- Measure  $x$ -dependence of nuclear effects
  - e.g.  $F_2$  and  $xF_3$ 
    - Are sea and valence quarks affected differently?
- Contribute to community efforts to extract the dependence of nuclear parton distribution functions on  $A$ ,  $x$  and  $Q^2$




# MINER $\nu$ A objectives

- Measure final state multiplicities as a function of A
  - Investigate pion absorption in the interaction nucleus
  - Observe final state interactions (FSI)
    - Improve models of  $\nu$ -A interaction
- Measure hadronic energy as a function of A
  - What is the A dependence of observed energy?
  - Also measure unobserved energy emission via neutrons
    - Improves neutrino energy resolution

# Neutrino flux

- [Understanding the neutrino flux](#)


1. Comparing  $\nu$  event rates when varying beamline configuration (target z-position and horn current). Each configuration samples different pion kinematics at the same neutrino energy (8 different configurations in all)



# Vertex Z Distribution

**Iron**-Rich Sample  
**Lead**-Rich Sample  
**Plastic** Sample

Events Selected as Nuclear Target Events



- Main contamination is first active module downstream of passive material
  - ALL events in passive target sample have a reconstructed vertex in this module